arXiv:1207.3707v2 [cond-mat.quant-gas] 22 Aug 2012

Higgs Excitation in Cold Atoms with Spin-orbit Coupling

Fei-Jie Huang, Qi-Hui Chen, and Wu-Ming Liu
National Laboratory for Condensed Matter Physics, Institute of Physics,
Chinese Academy of Sciences, Beijing 100190, China

We investigate the Higgs excitation in cold atoms induced by non-Abelian gauge potential. We demonstrate that when a non-Abelian gauge potential reduces to Abelian potential, only its Abelian part suits to construct spin-orbit coupling, and its non-Abelian part emerges as a Higgs excitation. The Higgs excitation induces the generation of mass of the non-Abelian gauge field, and it can affect the spin Hall currents which are emerged by the spin-orbit coupling. We also discuss the observation of these phenomena in real experiment.

PACS numbers: 03.75.Ss, 03.65.Vf, 67.85.Lm, 67.85.De

Introduction.— Spin-orbit (SO) coupling, which is the interaction between the spin and the momentum of a particle, relates to many effects in condensed physics, especially it is essential for realizing spin-Hall effect [1, 2] and topological insulators [3, 4] in solid systems. Recently, with realizing artificial gauge potentials [5], the cold atom systems, which have been proven to be perfect quantum emulators [6, 7], also can be allowed to simulate SO coupling [8–10]. This opens a new arena to explore various effects in cold atom systems [11–14]. The key to simulate SO coupling in cold atom systems is the generation of non-Abelian gauge potentials, which are synthesized from the interaction between the atoms and the lasers [5]. It is well known that the non-Abelian gauge potentials play a crucial role in understanding the fundamental interactions in particle physics. In condensed physics, the non-Abelian gauge potentials also appear in learning the mechanisms of high T_c superconductors [15] and graphene [16]. Generally, the non-Abelian gauge potentials in solid systems are concerned with spin. It is no doubt that, to emerge SO coupling, the minimal symmetry of the gauge potentials is required to be $SU(2)$.



FIG. 1. (Color online) (a) Atoms with Λ-level configuration interact with laser beams. $\mid g_1\rangle$ and $\mid g_2\rangle$ are the ground states, $\mid e\rangle$ is the excited state. Ω_1 and Ω_2 are the Rabi frequencies, Δ is a large detuning. The non-Abelian potential is arisen by two degenerate eigenstates which are induced by the large detunning. (b) The configuration of laser fields. The laser fields have the same wave numbers $\mid \mathbf{k} \mid = \mid \mathbf{k}' \mid$, there is an angle φ between them. Two inner laser fields (the blue arrows) are arranged to form Rabi frequency $\Omega_1 = \frac{1}{2}\Omega[\exp(i\mathbf{k}\cdot\mathbf{r}) + \exp(i\mathbf{k}'\cdot\mathbf{r})]$, other two lasers (the red arrows) are arranged to form $\Omega_2 = \frac{1}{2i}\Omega[\exp(i\mathbf{k}\cdot\mathbf{r}) - \exp(i\mathbf{k}'\cdot\mathbf{r})]$.

The generation mechanism of mass is one of the core issues for the whole physics, the richness of the world depends on the existence of mass particles. The mass generation mechanisms refer to the Higgs field. Various experiments have been designed to search the Higgs particle to confirm the Higgs field. Recently, the Large Hadron Collider (LHC) announced that they had found the suspected Higgs particle, and that may be the most breakthrough in particle physics in two decades [17, 18]. One of the advantage of introducing the Higgs field is that it can solve the mass generation mechanisms of the gauge field. Either through Higgs mechanism or through the symmetry reduction of gauge potential, the mass of gauge field can be obtained by the Higgs field. When referring to the reduction of $SU(2)$ gauge potential, if the reduction dose not satisfy a translation condition, the mass of gauge field will be generated by the Higgs excitation [19–23], which is induced by the Higgs field [24, 25]. Yet generally in particle physics, the meaningful reduction is to emerge a gauge charge, such as the 't Hooft monopole [26], which normally satisfies the translation condition and couples to gauge field only. In order to generate the mass of gauge field, a more broad charge should be taken into account. Therefore, the SO coupling is an option. In presence of SO coupling, the spin can be viewed as a charge which not only couples to the gauge field, but also to the orbit of the system.

In this Letter, we will investigate the Higgs excitation, as the generation of mass of the gauge field, by constructing the SO coupling in a cold atom system with non-Abelian gauge potential. In cold atom systems, for their unprecedented controllability, the reduction of symmetry would own its unique features. It is well known that the SO coupling appears in $U(1)$ system, so to construct SO coupling, one should reduce the $SU(2)$ gauge potential to $U(1)$ potential. On the other hand, the spin can be viewed as a charge which not only couples to the gauge field but also to the orbit. Thus we can expect that through the reduction of $SU(2)$ gauge potential, an excitation may appear in constructing of SO coupling, and the mass of the gauge field can be generated. We



FIG. 2. (Color online) The sphere surface of gauge potential $\mathcal{A}_0$ with $SU(2)$ symmetry. (a) At the initial time $t = 0$, the base vectors of the gauge potential $\sigma_x(0)$, $\sigma_y(0)$ and $\sigma_z(0)$ point to certain directions. The direction vector (the red arrow) $\sigma(x)$ points to A. (b) After a time t, the base vectors change to $\sigma_x(t)$, $\sigma_y(t)$ and $\sigma_z(t)$ directions. The direction vector $\sigma(x)$ changes along with the base vectors and points to B. If the path AB satisfies the equation of motion of spin, the $SU(2)$ potential $\mathcal{A}_0$ will induce the appearance of SO coupling. If not, then excepting for SO coupling, the gauge covariant potential $\mathcal{B}$ also appears. The potential $\mathcal{B}$ will induce the Higgs excitation.

also discuss the signatures of these phenomena in real experiment.

The spin-orbit coupling in non-Abelian gauge potential.— We consider a cold atom system with Λ-type level as shown in Fig. 1a. The lasers are arranged to couple two ground states $\mid g_1\rangle$ and $\mid g_2\rangle$ to an excited state $\mid e\rangle$. The Rabi frequencies are taken as $\Omega_1 = \frac{1}{2}\Omega[\exp(i\mathbf{k}\cdot\mathbf{r}) + \exp(i\mathbf{k}'\cdot\mathbf{r})]$ and $\Omega_2 = \frac{1}{2i}\Omega[\exp(i\mathbf{k}\cdot\mathbf{r}) - \exp(i\mathbf{k}'\cdot\mathbf{r})]$, in which $\mathbf{k}$ and $\mathbf{k}'$ are the wave vectors of lasers, $\mathbf{k}' = e^{i\varphi}\mathbf{k}$, $\mathbf{r}$ is the position vector, φ is the angle between the lasers as shown in Fig. 1b. The total Rabi frequency is given by $\Omega = (|\Omega_1|^2 + |\Omega_2|^2)^{1/2}$.

Defining the total wave vector $\mathbf{Q} = \mathbf{k} + \mathbf{k}'$ and the relative wave vector $\mathbf{q} = \mathbf{k} - \mathbf{k}'$, the Rabi frequencies can be redefined as $\Omega_1 = \Omega\cos\theta\exp(i\phi_1)$ and $\Omega_2 = \Omega\sin\theta\exp(i\phi_2)$, in which the parameters are $\theta = \frac{1}{2}\mathbf{q}\cdot\mathbf{r}$, $\phi_1 = \phi_2 = \frac{1}{2}\mathbf{Q}\cdot\mathbf{r}$. The Hamiltonian of the system reads $H = H_k + H_I + V_{trap}$, in which $H_k = p^2/2m$ is the kinetic energy, p is the momentum of single cold atom, m is the atomic mass, V_{trap} is a trapping potential. The interacting Hamiltonian is given by $H_I = 2\Delta \mid e\rangle\langle e \mid + (\Omega_1 \mid e\rangle\langle g_1 \mid + \Omega_2 \mid e\rangle\langle g_2 \mid + h.c.)$, where Δ is the detuning.

The eigenstates of the interacting Hamiltonian can be written as $\mid\chi_{D1}\rangle = -\sin\theta \mid g_1\rangle + \cos\theta \mid g_2\rangle$, $\mid\chi_{D2}\rangle = \cos\delta\cos\theta \mid g_1\rangle + \cos\delta\sin\theta \mid g_2\rangle - \sin\delta e^{-i\phi_1} \mid e\rangle$, $\mid\chi_3\rangle = \sin\delta\cos\theta \mid g_1\rangle + \sin\delta\sin\theta \mid g_2\rangle + \cos\delta e^{-i\phi_1} \mid e\rangle$, where $\tan\delta = \frac{1}{\Omega}[(\Delta^2+\Omega^2)^{1/2} - \Delta]$, the eigenvalues are $E_{D1} = 0$, $E_{D2,3} = [\Delta \mp (\Delta^2 + \Omega^2)^{1/2}]$.

For large detuning, $\tan\delta << 1$, the eigenvalue $E_{D2} \to 0$, and the two lower internal eigenstates will give rise an effective $SU(2)$ gauge potential $\mathbf{A} = \frac{1}{2}\mathbf{q}\sigma_y + \frac{1}{4}\delta^2\mathbf{Q}\sigma_z$. Redefining this effective $SU(2)$ gauge potential to a Lorentz scalar potential $\mathcal{A}_0 = [\gamma, \mathbf{A}]_+$, where dimensionless parameter $\gamma = \mathbf{p}/p$, $[,]_+$ is an anti-commutator. Then neglecting the trapping potential and the constant terms, the effective Hamiltonian of the system can be written as

$$H = H_k + g\mathcal{A}_0 + V, \tag{1}$$

where $H_k = \eta p$ is the kinetic energy, $g = \eta$ is an effective charge, $\eta = p/2m$ is a dimensionless strength factor referring to the non-relativistic approximation (assuming the light velocity $c = 1$). $V = [\frac{1}{16m}\delta^2(q^2 - (1+\delta^2)Q^2) + \delta\frac{\Omega}{2}]\sigma_z = M_0\sigma_z$ is a scalar potential which originates from constructing of $\mathbf{A}$.

It is well known that the SO coupling turns up in a $U(1)$ gauge potential, so in order to construct SO coupling, let's reduce the $SU(2)$ gauge potential to $U(1)$ symmetry. Firstly, we denote the unit direction $\mathbf{n} = (n_x, n_y, n_z)$ along the direction of magnetic field, and define the internal $SU(2)$ space direction vector as $\sigma(x) = n_i\sigma_i(t)\hat{i}, x = (t, \mathbf{x}), i = x, y, z$, which means that we have chosen the same directions for the external and the internal space. Then along the direction vector $\sigma(x)$, $\mathcal{A}_0$ will decompose to two gauge potentials [24, 27], which can be read as $\mathcal{A}_0 = \mathcal{A} + \mathcal{B}$, with

$$\begin{aligned}\mathcal{A} &= (\sigma(x)\cdot\mathcal{A}_0)\sigma(x) + [\partial_0\sigma(x), \sigma(x)],\\ \mathcal{B} &= [\sigma(x), \nabla_0\sigma(x)],\end{aligned} \tag{2}$$

where $\nabla_0 = \partial_0 + [\mathcal{A}_0,]$ represents the time component of covariant differentiation, $[,]$ denotes the commutator. According to differential geometry theorem [28], when $\mathcal{A}_0$ reduces to $U(1)$ potential, there should be $\nabla_0\sigma(x) = 0$. Therefore, if $\mathcal{A}_0$ is a $U(1)$ potential, $\mathcal{A}_0 = \mathcal{A}$. We view $\mathcal{A}$ is the Abelian part of $\mathcal{A}_0$, and the potential $\mathcal{B}$ is the non-Abelian part.

Now let's discuss the physical meaning of the constraint condition

$$\nabla_0\sigma(x) = 0. \tag{3}$$

This condition actually is a translation condition. Taking the Planck constant $\hbar = 1$, $\partial_0 = i\,\partial_t$ corresponds to the Hamiltonian operator of the system. Then neglecting the scalar potential V and denoting the effective charge $g = 1$, we obtain $[\sigma(x), H] = [\sigma(x), \mathcal{A}_0]$, and the constraint condition equals to $i\partial_t\sigma(x) = [\sigma(x), H]$. This equation just is the equation of motion of spin. Thus the constraint condition in fact corresponds to the equation of motion of spin, and the direction vector $\sigma(x)$ corresponds to the spin operator of the system. Therefore, we find that the constraint condition is the relationship between the spin and the $SU(2)$ gauge potential, as shown in Fig. 2.

We conclude that, only when the non-Abelian gauge potential $\mathcal{A}_0$ satisfies the constraint condition $\nabla_0\sigma(x) = 0$, $\mathcal{A}_0$ can be used to generate the SO coupling. If the constraint condition can not be satisfied, the gauge potential $\mathcal{B}$ in $\mathcal{A}_0$ will can't be canceled. In this case, $\mathcal{A}_0$ will not



FIG. 3. (Color online) (a) The trajectories of cold atoms contributed by the Higgs excitation. The red solid line and the blue dash line correspond to the velocities $v = 3 \times 10^{-3} \mu m/s$ and $v = 5 \times 10^{-3} \mu m/s$. It is shown that the contribution includes two opposite trajectories, one along the $\mathbf{r}$ direction, the other is inverse. (b) The dispersion of coupling energy between the Higgs excitation and the atoms, $\varepsilon_B = (\epsilon p + \eta' M_B)\sigma_z$, in which p is the momentum of the single atom, M_B is the excited mass. The red solid line and the blue dash line correspond to the coupling strength parameters $\epsilon = 4m^{-1} \times 10^{-2}$ and $\epsilon = 6m^{-1} \times 10^{-2}$, m is the atomic mass. The unit of the energy ε_B is the recoil energy $\varepsilon_R = k^2/(2m)$, k is the wave number of the lasers. Due to the presence of the excited mass M_B, the dispersion is not vanish at zero momentum.

suit to construct SO coupling, because if the equation of motion of spin is absent, the SO coupling is also absent in the system. Therefore, the equation $\mathcal{A}_0 = \mathcal{A}$ sets the $SU(2)$ gauge potentials which are drawn up to construct SO coupling.

We now catch up on the scalar potential V into the effective Hamiltonian to check whether $\mathcal{A}_0$ satisfies the constraint condition $\nabla_0 \sigma(x) = 0$ or not. The constraint condition can be rewritten as $\nabla_0 \sigma(x) = [\sigma(x), H] + [\mathcal{A}_0, \sigma(x)]$. On this occasion, because the scalar potential is a non-Abelian potential, we find $\nabla_0 \sigma(x) \neq 0$. It indicates that the potential $\mathcal{B}$ will exist according to the Eq. (2). In this case, $\mathcal{A}_0$ does not suit to construct SO coupling, thus the SO coupling term of the cold atoms system should be read as $H_{SO} = g\mathcal{A}$, with its explicit expression

$$H_{SO} = \lambda \sigma_y + \nu \sigma_z. \tag{4}$$

The coupling strength factors λ and ν are $\lambda = \frac{1}{m}[\frac{1}{2}\mathbf{q} \cdot \mathbf{p} + \delta^3 \Omega p (\mathbf{Q} \cdot \mathbf{p} / \mathbf{q} \cdot \mathbf{p})]$, $\nu = \frac{1}{m}(-2\delta\Omega p + 2^{-2}\delta^2 \mathbf{Q} \cdot \mathbf{p}) + \frac{1}{4m^2}\delta^2(Q^2 - q^2)p$, in which the direction of $\sigma(x)$ is chosen to consist with the initial direction of $\mathcal{A}_0$. The Eq. (4) is a generalized expression of SO coupling, it shows that other than only one component, all components of momentum can participate in the SO coupling.

Higgs excitation in cold atom system.— Owning to the non-Abelian potential $\mathcal{A}_0$ can not satisfy the constraint condition, after constructing SO coupling, there will be a surplus term left in the Hamiltonian. This term can be expressed as $H_{k,B} = g\mathcal{B}$, with $\mathcal{B} = -\frac{\mathbf{Q} \cdot \mathbf{p}}{\mathbf{q} \cdot \mathbf{p}} \delta^3 \Omega \sigma_y + [2\delta\Omega + \frac{1}{4m}\delta^2(Q^2 - q^2)]\sigma_z$. The potential $\mathcal{B}$ is a gauge covariant potential, it can own a mass term $M_B^2 Tr[\mathcal{B} \cdot \mathcal{B}]$. By writing down the action of non-Abelian gauge field, the mass reads $M_B = (1 + (\mathbf{Q} \cdot \mathbf{p} / 2\mathbf{q} \cdot \mathbf{p})^2 \delta^4)^{-\frac{1}{2}} M_0$. Thus the effective Hamiltonian of the system can be expressed as



FIG. 4. (Color online) The relationship between the particle number N and the trap frequency ω. ε_F is the Fermi energy. The red solid line and the green dash line correspond to the ratios between the excited mass M_B and a characteristic mass m^*, $M_B/m^* = 2.2$ and $M_B/m^* = 2.8$ respectively. $m^* = q^2/2m$ is a characteristic mass of the system, in which q is the relative momentum of the laser fields.

$$H = H_k + H_{SO} + H_B, \tag{5}$$

with

$$H_B = H_{k,B} + \eta' M_B \sigma_z. \tag{6}$$

It is clear that an excitation appears in the Eq. (6). $H_{k,B} = g\mathcal{B}$ describes the coupling between the kinetic energy of the excitation and the atoms. The scalar potential $V = \eta' M_B \sigma_z$ in fact describes the coupling between the mass of the excitation and the atoms, in which $\eta' = (1 + (\mathbf{Q} \cdot \mathbf{p} / 2\mathbf{q} \cdot \mathbf{p})^2 \delta^4)^{\frac{1}{2}}$ is a dimensionless coupling factor.

Let's discuss the origin of the excitation. As shown in Fig. 2, the original direction of the direction vector $\sigma(x)$ presents $SU(2)$ symmetry in the internal space, its direction can be chosen in freely. Nevertheless, when $\mathcal{A}_0$ is used to construct SO coupling, the direction of $\sigma(x)$ is constrained to the constraint condition, and the initial symmetry is broken. Yet in general, the whole $\mathcal{A}_0$ can not used to construct SO coupling potential, so there is a trend to restore the initial symmetry. The potential $\mathcal{B}$ just plays as the excitation which excites to restore the initial symmetry. We treat this excitation as a Higgs excitation, in which $\sigma(x)$ plays as the Higgs field [24, 25]. Due to the excitation, the mass of the gauge field $\mathcal{A}_0$ is generated.

The trajectories of cold atoms contributed by the excitation can be obtained by the equation of motion $\dot{\mathbf{r}} = -i[\mathbf{r}, H_B]$. As shown in Fig. 3a, the contribution includes two opposite trajectories. The Fig. 3b shows the dispersion of H_B. The dispersion is linear, and the energy is not vanish at zero momentum due to the presence of the excited mass.

Next, let's discuss the impact of the excitation to the cold atoms system. A two-dimensional harmonic potential $\frac{1}{2}m\omega^2(y^2+z^2)$ is chosen to trap the cold atoms. The relationship between the particle number and the trap frequency under the impact of the excited mass can be obtained by solving the equation $N=\int d\mathbf{r}n(r,t=0,T=0)$, in which N is the atomic number, $n(r,t,T)=\frac{1}{(2\pi)^2}\int d\mathbf{p}[f_+(p,r,t,T)+f_-(p,r,t,T)]$ is the density of the cold atoms at the time t, T is the temperature, $f_{\sigma_z}=f_\pm(p,r,t,T)=[e^{\beta(H_\pm(p,r,t)-\mu)}+1]^{-1}$ are the spin-depending Fermi distributions with $\beta=1/k_BT$. The relationship is shown in Fig. 4.

We now discuss the spin Hall currents of the system. To generate spin Hall current, the wave vectors of the lasers is chosen to $k_x=k'_x=0$, the internal space rotates $\pi/2$ around the x direction. In this case, the SO coupling reads $H_{SO}=\lambda\sigma_z$. This term describes the spin Hall currents in which the spin polarizes to z direction and the currents move along y direction. The spin Hall currents can be expressed as

$$J^y_{\sigma_z}=\frac{1}{(2\pi)^2}\int d\mathbf{p}f_{\sigma_z}(p)j^y_{\sigma_z}, \tag{7}$$

where $j^y_{\sigma_z}=\langle\hat{j}^y_{\sigma_z}\rangle$ is the single particle current. The spin current operator $\hat{j}^y_{\sigma_z}=\frac{1}{4}[\sigma_z,v_y]_+$, $v_y=-i[y,H]$ is the velocity along y direction. The impact of the excitation to the spin Hall currents can be shown as Fig. 5a. It can be seen that the spin down current is suppressed by the increase of the excited mass, while the spin up is slightly grown. The evolution of the atomic density $n(r,t)$ is shown in Fig. 5b.

Experimental signatures of Higgs excitation.— Let's discuss the observation of the Higgs excitation by detecting the spin Hall currents. We can choose ^6Li atoms for a Λ-type level system, the particle number is about 10^4, the bias magnetic field is about $10G$ [10], and a $2\pi\times10^2Hz$ harmonic potential is used to trap the atoms. The configuration of four laser fields is shown in Fig. 1b. The wave number of the lasers can be taken as $2\pi\times1.0(\mu m)^{-1}$ [10]. For a large detuning, the Rabi frequency and the detuning is required to satisfy $\Omega^2/\Delta\sim10^6Hz$. When the laser fields are turned on, a non-Abelian gauge potential is applied to the ^6Li atoms. By tuning the angle φ between the lasers, different masses of excitation can be obtained.

To detect the spin current, we firstly initialize the atoms in $\mid g_1\rangle$ state. Then a Raman pulse is applied between the states $\mid g_1\rangle$ and $\mid g_2\rangle$ to transfer the atoms to spin up state $\mid\chi_{D1}\rangle$. The Rabi frequency of the Raman pulse is required to match the spatial variation of the $\mid\chi_{D1}\rangle$ state. Turning on the lasers, the cold atoms will experience the SO coupling, and couple to the excitation. After a time t, we turn off the lasers and transfer the atoms from $\mid\chi_{D1}\rangle$ state back to the initial state $\mid g_1\rangle$ by applied a reversal Raman pulse. In this case, by using the time of flight measurement, we can determine the spin up current of the system. The measurement of the spin down current (corresponding to the $\mid\chi_{D2}\rangle$ state) also can be detected in the same manner [29].



FIG. 5. (Color online) (a) The relationship between the excited mass M_B and the spin Hall currents $J^y_{\sigma_z}$. The red (blue) solid line is correspondence to the spin up (down). The unit of $J^y_{\sigma_z}$ is $\frac{1}{2\pi}\times10cm/s$. We choose ^6Li atoms and the wave number of the lasers is $2\pi\times1.0(\mu m)^{-1}$, the particle number is about 10^4. It can be seen that the spin down current is suppressed by the increase of the excited mass, while the spin up is slightly grown. (b) The evolution of the atomic density from the time $t=0ms$ to $t=4ms$ and $t=9ms$. The unit of y and z directions is μm. The temperature is taken as $T=0.4T_F$, T_F is the Fermi temperature. The up and down figures correspond to the ratios $M_B/m^*=2.2,\ 2.8$. The red and blue parts in each figure denote the spin up and spin down currents. The figures show that the variation trends of spin currents are the same as Fig. 5a.

Conclusion.— In summary, by constructing SO coupling from non-Abelian gauge potential, we have investigated the Higgs excitation, which leads to the generation of mass of the gauge field, in cold atom system. We argue that when a non-Abelian gauge potential reduces to Abelian potential, only its Abelian part suits to construct SO coupling, and its non-Abelian part emerges as a Higgs excitation. The excitation can possess mass, and it can affect the spin Hall currents. We also discuss how to observe the excitation through the detection of spin Hall currents in experiment. We expect that the emergence of the Higgs excitation in cold atoms can help to understand the generation mechanics of mass, which is interesting from the fundamental interaction theories to condensed matter physics.

This work was supported by NKBRSFC under grants Nos. 2011CB921502, 2012CB821305, 2009CB930701, 2010CB922904, the NSFC under grants Nos. 10934010, 60978019, and NSFC-RGC under grants Nos. 11061160490 and 1386-N-HKU748/10.

[1] S. Murakami, N. Nagaosa, and S. -C. Zhang, Science 301, 1348 (2003).
[2] J. Sinova, D. Culcer, Q. Niu, N. A. Sinitsyn, T. Jungwirth, and A. H. MacDonald, Phys. Rev. Lett. 92, 126603 (2004).
[3] M. Z. Hasan and C. L. Kane, Rev. Mod. Phys. 82, 3045 (2010).
[4] X. -L. Qi, S. -C. Zhang, Rev. Mod. Phys. 83, 1057 (2011).
[5] J. Dalibard, F. Gerbier, G. Juzeliūnas, and P. Öhberg, Rev. Mod. Phys. 83, 1523 (2011).
[6] I. Buluta and F. Nori, Science 326, 108 (2009).
[7] I. Bloch, J. Dalibard and S. Nascimbéne, Nature Phys. 8, 267 (2012).
[8] Y. -J. Lin, K. Jiménez-Garcia, J. V. Prto, and I. B. Spielman, Narure 471, 83 (2011).
[9] P. Wang, Z. Yu, Z. Fu, J. Miao, L. Huang, S. Chai, H. Zhai, and J. Zhang, arXiv:1204.1887.
[10] L. W. Cheuk, A. T. Sommer, Z. Hadzibabic, T. Yefsah, W. S. Bakr, and M. W. Zwierlein, arXiv:1205.3483.
[11] J. Ruseckas, G. Juzeliunas, P. Öhberg, and M. Fleischhauer, Phys. Rev. Lett. 95, 010404 (2005).
[12] S. -L. Zhu, H. Fu, C. -J. Wu, S. -C. Zhang, and L. -M. Duan, Phys. Rev. Lett. 97, 240401 (2006).
[13] X. -J. Liu, M. F. Borunda, X. Liu, and J. Sinova, Phys. Rev. Lett. 102, 046402 (2009).
[14] R. Liao, Y. Yi-Xiang, and W. -M. Liu, Phys. Rev. Lett. 108, 080406 (2012).
[15] P. A. Lee, N. Nagaosa, X. G. Wen, Rev. Mod. Phys. 78, 17 (2006).
[16] M. A. H. Vozmedianoa, M. I. Katsnelsonb, F. Guinea, Phys. Rep. 496, 109 (2010).
[17] J. Incandela, CMS talk at *Latest update in the search for the Higgs boson* at CERN, July4, 2012, https:// cms-docdb.cern.ch/cgi-bin/PublicDocDB//ShowDocument?docid=6125.
[18] F. Gianotti, ATLAS talk at *Latest update in the search for the Higgs boson* at CERN, July4, 2012, https:// cms-docdb.cern.ch/cgi-bin/PublicDocDB//ShowDocument?docid=6126.
[19] S. D. Huber, B. Theiler, E. Altman, and G. Blatter, Phys. Rev. Lett. 100, 050404 (2008).
[20] D. Podolsky, A. Auerbach, and D. P. Arovas, Phys. Rev. B 84, 174522 (2011).
[21] F. Cooper, C. C. Chien, B. Mihaila, J. F. Dawson, and E. Timmermans, Phys. Rev. A 85, 023631 (2012).
[22] L. Pollet and N. Prokof'ev, Phys. Rev. Lett. 109, 010401 (2012).
[23] M. Endres, T. Fukuhara, D. Pekker, M. Cheneau, P. Schauβ, C. Gross, E. Demler, S. Kuhr, and I. Bloch, Narure 487, 454 (2012).
[24] Y. S. Duan and M. L. Ge, Sinica Sci. 11, 1072 (1979).
[25] L. Faddeev and A. J. Niemi, Nucl. Phys. B 776, 38 (2006).
[26] G. 't Hooft, Nucl. Phys. B 79, 276 (1974).
[27] Y. M. Cho, Phys. Rev. D 21, 1080 (1980).
[28] S. Kobayashi and K. Nomizu, *Foundations of Differential Geometry*, vol.1 and vol.2 (Wiley-Interscience, New York, 1996).
[29] T. D. Stanescu, C. Zhang and V. Galitski, Phys. Rev. Lett. 99, 110403 (2007).